

Discretization on the Cosmic Scale Inspired from the Old Quantum Mechanics

A. G. Agnese* R. Festa[†]

To appear in: *Proceedings of the Beer Sheva Workshop on Modern Modified
Theories of Gravitation and Cosmology* (1997)

Abstract

The Old Quantum Mechanics action discretization rules for periodic motions on the atomic scale (Bohr-Sommerfeld) have been suitably modified in order to take into account the gravitational field instead of the electrostatic one. The new rules are used to calculate a few mechanical quantities pertinent to the periodic motions of celestial objects, and several values are obtained which result in reasonable agreement with the corresponding experimental data.

A gravitational dimensionless structure constant α_g is determined, using the data relative to the solar system, which allows to quantitatively account for phenomena on a much wider scale. In particular, some information is acquired about the recently discovered extrasolar planetary systems and about the general empirical law which connects the spin of a celestial body with the square of its mass.

Though no general theory is explicitly proposed in order to explain the obtained numerical coincidences, Nelson's stochastic mechanics and/or chaotic dynamics seem to be the best candidates for this rôle.

*Università di Genova, Dipartimento di Fisica, and INFN, Sezione di Genova, Via Dodecaneso 33, 16146 Genova (Italy). E-Mail: agnese@genova.infn.it .

[†]Università di Genova, Dipartimento di Fisica, and INFN, Unità di Genova, Via Dodecaneso 33, 16146 Genova (Italy). E-Mail: festa@genova.infn.it .

I. Introduction

In a recent work [?] we suggested that the periodic motions of celestial bodies can be described, approximately enough, starting from suitably tuned action discretization rules (scaling up of the Bohr-Sommerfeld laws). In fact, if the Planck's constant is expressed using the relationship

$$h = 2\pi \frac{e^2}{\alpha_e c} ,$$

where $\alpha_e \simeq 1/137$ is the dimensionless fine structure constant, the Bohr-Sommerfeld rules can be written in the form

$$\oint p_j dq_j = n_j 2\pi \frac{e^2}{\alpha_e c} .$$

One can imagine that, if the electron charge was known with higher precision in Planck's time, he would have introduced α_e instead of h as a new constant.

The form of the previous equation explicitly shows its 'atomic electrostatic' purpose. On the other hand, since both the (atomic) electrostatic and the gravitational force field follow the r^{-2} law and only differ in their 'sources':

$$\begin{aligned} e^2 &= \text{proton charge times electron charge in the H atom} \\ GMm &= \text{mass m in the gravitational field of a source of mass M} , \end{aligned}$$

we will conjecture that the Old Quantum Mechanics rules hold if one performs the substitution

$$\frac{e^2}{\alpha_e} \longrightarrow \frac{GMm}{\alpha_g} ,$$

where α_g is a (new) dimensionless 'gravitational' structure constant to be determined.

Introducing the gravitational potential energy

$$V = -\frac{GMm}{r} ,$$

it follows by simple calculations that the major semi-axes of the (elliptical) orbits are given by

$$a_n = n^2 a_1 ,$$

where $n = 1, 2, \dots$ is the principal number and

$$a_1 = \frac{GM}{\alpha_g^2 c^2}$$

is the ‘Bohr radius’ of the M gravitational source.

Other immediate consequences concern the minimum and maximum orbital distances q_{nl} and Q_{nl} and the corresponding maximum and minimum speeds V_{nl} and v_{nl} :

$$\begin{cases} q_{n,l} &= a_n(1 - \varepsilon) & ; & V_{n,l} &= (1 + \varepsilon) \frac{\alpha_g c}{l} \\ Q_{n,l} &= a_n(1 + \varepsilon) & ; & v_{n,l} &= (1 - \varepsilon) \frac{\alpha_g c}{l} \end{cases} ,$$

where

$$\varepsilon = \sqrt{1 - \frac{l^2}{n^2}}$$

is the elliptical eccentricity, and, as customary, the azimuthal quantum number l can take the values $1, \dots, n$. The angular momentum is given by

$$j_{n,l} = l \frac{GMm}{\alpha_g c} .$$

In the case of circular orbits ($l = n$) one has the constant speed

$$v_n = \frac{\alpha_g c}{n} ,$$

the corresponding period

$$P_n = 2\pi \frac{GM}{\alpha_g^3 c^3} n^3 ,$$

and the angular momentum

$$j_n = \frac{GMm}{\alpha_g c} n .$$

Thus, the universal constant

$$v_* = \alpha_g c$$

turns out to be the maximum speed for a gravitationally bounded orbitating point mass. Correspondingly,

$$P_* = 2\pi \frac{GM}{\alpha_g^3 c^3} ,$$

which depends on the strength M of the gravitational source, turns out to be its minimum period, and

$$j_* = \frac{GMm}{\alpha_g c}$$

its minimum angular momentum.

II. A ‘domestic’ test case: orbital motions around the Sun

As a first test case we can apply the previous formulae to the motion of the solar planets. In this case we use the following relationships

$$\left\{ \begin{array}{ll} \text{Major semi-axes :} & a_n = n^2 a_1^\odot , \quad \text{where } a_1^\odot = \frac{GM^\odot}{\alpha_g^2 c^2} \\ \text{Periods :} & P_n = n^3 P_1^\odot , \quad \text{where } P_1^\odot = \frac{2\pi GM^\odot}{\alpha_g^3 c^3} \\ \text{Mean speeds :} & v_n = \frac{\alpha_g c}{n} \end{array} \right.$$

where $M^\odot$ indicates the mass of the Sun.

In order to determine α_g , we previously need to select the quantum principal numbers for the various planets. Note that, when a given number for a given planet is chosen, the numbers for all the other planets turn out to be determined by their observed mean speeds. Of course, the results are significant if the concerned quantum principal numbers are small, and if no satisfactory agreement exists for semi-integer n values.

One can easily verify that the only consistent choice lies in assigning to Mercury the principal number 3. Using the corresponding known speed one obtains the initial guess for the gravitational structure constant

$$\alpha_g = 3 \frac{v_{Mercury}}{c} ,$$

and the following quantum number selection: Mercury→3, Venus→4, Earth→5, Mars→6, Jupiter→11, Saturn→15, Uranus→21, Neptune→26, Pluto→30.

On this basis, one can use the relationship $v_n = \alpha_g c/n$ to refine the estimate of α_g through a general least square fit.

In this way, the following results can be obtained:

$$1/\alpha_g = 2086 \pm 14 ,$$

$$v_* = 143.7 \pm 1.0 \text{ km/s} ,$$

$$a_1^\odot = 0.04297 \text{ a.u.} ,$$

$$P_1^\odot = 3.269 \text{ d} ,$$

and

Planet	n	$v_{obs} \text{ (km/s)}$	$v_{calc} \text{ (km/s)}$
Mercury	3	47.87	47.90
Venus	4	35.02	35.92
Earth	5	29.78	28.74
Mars	6	24.13	23.95
Jupiter	11	13.06	13.06
Saturn	15	9.64	9.58
Uranus	21	6.80	6.84
Neptune	26	5.43	5.53
Pluto	30	4.74	4.79

In Fig. 1 the $v_n = \alpha_g c/n$ and $a_n = n^2 a_1$ relationships are graphically compared with the actual orbital mean speeds and the actual major semi-axes of the planets of the Sun. One can see that the solar planetary motions are reasonably accounted by our model of ‘gravitational atom’, even for small principal numbers.

As the azimuthal numbers are concerned, considering one of the two planets with valuable eccentricity, i.e. Pluto ($n = 30$), and using the eccentricity formula with $l = 29$, one obtains $\varepsilon_{calc} = 0.256$, in fair agreement with the observed value $\varepsilon_{obs} \simeq 0.25$. Unfortunately, no allowable calculated value fits satisfactorily the eccentricity of Mercury (whose motion shows on the other hand the well known anomalies).

Very recently [?] a new object has been discovered, named 1996 TL_{66} , which can be assumed to adequately represent a number of bodies located between the Kuiper belt (beyond the Neptune orbit) and the Oort cloud (which reaches about 50,000 a.u. from the Sun). The orbit of this object,

the brightest trans-neptunian body but Pluto and Charon, shows the very large eccentricity $\varepsilon = 0.58$ and a major semiaxis $a_n = 83.7 a.u.$.

Using our mechanical model, one gets the (n, a_n) pairs $(43, 79.4 a.u.)$, $(44, 83.2 a.u.)$ and $(45, 87.0 a.u.)$: thus, the principal number $n = 44$ seems to be adequate to this ‘new planet’. ‘Scanning now the allowable principal-azimuthal number pairs we obtain $(44, 35) \rightarrow \varepsilon = 0.61$, $(44, 36) \rightarrow \varepsilon = 0.58$ and $(44, 37) \rightarrow \varepsilon = 0.54$, so that the pair $(n, l) = (44, 36)$ can be assigned to $1996TL_{66}$. Nevertheless, we explicitly agree on the fact that results based on such large quantum numbers can not contribute to confirm or disprove our model.

III. Orbital motions around the planets

On the basis of the assumed universality of the constant α_g one can use, in evaluating the ‘Bohr radius’ of each planet, the formula

$$r_1^{(planet)} = a_1^\odot \frac{M^{(planet)}}{M^\odot},$$

Through $r_1^{(planet)}$ one can easily calculate the allowed orbits relative to each planet. In particular, one can determine the principal number n_f which corresponds to the first actual ‘free’ orbit (i.e., to the allowed orbit immediately out of the planet body). In the following table we show some interesting results (all the lengths are reported in km, and the general relative uncertainty of the calculated radii is about 1%). In the last two columns we report the greatest ‘not-free’ calculated orbital radius and the observed equatorial radius of the concerned body. Note that, in estimating n_{free} , the well known internal stability Roche limit [?] has not been taken into account: some underestimation could result.

Considering the principal number of the first ‘free’ orbit, one can note that that for the internal planets (and for Pluto) these orbits correspond to relatively high (≥ 19) principal numbers. The opposite, i.e. small (≤ 10) ‘free’ quantum numbers, is verified for the big external planets.

Body	Bohr radius	n_{free}	$r_{n_{free}}$	$r_{n_{free}-1}$	Eq. radius
Sun	$6.50 \cdot 10^6$	1	$(6.50 \pm 0.06) \cdot 10^6$		$6.96 \cdot 10^5$
Mercury	1.08	48	2488	2386	2439
Venus	15.86	20	6344	5725	6051
Earth	19.45	19	7021	6302	6378
Mars	2.09	41	3513	3344	3397
Jupiter	6190	4	99040	55710	71492
Saturn	1853	6	66708	46325	60268
Uranus	282	10	28200	22842	25559
Neptune	333	9	26918	21312	24764
Pluto	$4400 \cdot 10^{-4}$	160	1126	1112	1123

If the orbits of the natural satellites of the planets are considered one really obtains ambiguous results. For instance, only three of the four greater Jupiter satellites confirm our rule. This fact might denote some failure of our model or, on the contrary, that the fourth satellite does not originate from the same Jupiter-material as its companions. A particular case is that of the Earth-Moon pair: we will consider this case later.

As the Saturn rings are concerned, one obtains the following results, where the unit length is $10^3 km$, and IE , OE indicate respectively the internal and the external edge of the considered ring:

Rings	D_{IE}	C_{IE}	B_{IE}	Cassini	A_{IE}	F	G	E_{IE}	E_{OE}
r_{obs}	67.0	74.5	92.0	119.8	122.2	140.4	170	180	480
n	6		7		8	9		10	16
r_{calc}	66.7		90.8		118.6	150.1		185.3	474.4

The uncertainties of the calculated values are all about 1%. The calculated inner edge positions of most rings are in fair agreement with the corresponding experimental values. Note that the boundary of the C ring is presumably determined by the interactions between the charged dust particles and the planetary magnetic field, so that a purely gravitational theory does not account for this fact. Note also the remarkable coincidence between the calculated and the observed values of the boundaries of the broad rarefied E ring.

IV. Extrasolar planets

During the last two years a few extrasolar planetary systems have been discovered on the basis of the detected periodical anomalies in the motions of their stars. Unfortunately, the mass of most the concerned stars are at present estimated with some inaccuracy. Almost all the discovered planets, with masses ranging between one half and ten times the Jupiter mass, turn out to move somewhat near to the star, their guessed major semi-axes ranging from about 0.05 *a.u.* and 1 *a.u.* (only three planets show a mean distance from the central star greater than 1 *a.u.* and less than 2 *a.u.*).

In order to test our scheme, we have considered only the 19 extra-solar planets which have been confirmed (at present) [?] and two more planetary systems (Proxima Centauri and Barnard's) whose star masses are known with sufficient accuracy. The star masses of the other 19 confirmed planetary systems have been guessed from the star type information using the values M_{st} reported in literature ([?]). Then, the major semi-axes a_{st} of the planetary orbits have been calculated from the corresponding measured periods. Unfortunately, the starting point (i.e., the values of the star masses) was somewhat inaccurate.

Thus, an alternative path has been devised: starting from the measured planetary periods, and using the relationship

$$M_n = M_{\odot} \frac{P}{P_{\odot}} \frac{1}{n^3} ,$$

we scanned the various hypothetical star masses $M_1, M_2, \dots, M_n$ which would correspond to the various planet principal numbers $1, 2, \dots, n$. Then, the mass value M_j nearest to the guessed star-type mass M_{st} was singled out: thus, the the appropriate principal number j was determined , and this in turn allowed an appropriate estimate $a_j^{(est)}$ of the major semi-axis. In the following table, in order to put in evidence the ambiguity level in choosing the ‘most likely’ star mass, we report, together the chosen mass M_j , its neighbouring hypothetical values M_{j-1} and M_{j+1} .

We explicitly note that for most of the considered planetary systems the choice of the appropriate M_j is unequivocal enough.

For the last two planetary systems, the planet period was astrometrically determined. In both cases, the chosen M_j value turns out to fit exactly the measured one: this fact seems to support the correctness of our estimation method.

Star	Type	M_{st}	$P^{(obs)}$	a_{st}	M_{j-1}	M_j	M_{j+1}	n	$a_j^{(est)}$
51 Peg	G2IVA	1.05	4.229	0.050	-	1.30	0.16	1	0.056
ups Androm.	F7V	1.20	4.611	0.057	-	1.42	0.18	1	0.061
55 Cancer	G8V	0.90	14.648	0.110	4.50	0.56	0.17	2	0.097
rho CrB	G0V-G2V	1.05	39.645	0.230	12.18	1.52	0.45	2	0.261
16 Cyg B	G2.5V	1.05	804.000	1.720	1.98	1.14	0.72	6	1.766
47 Uma	G0V	1.05	1088.445	2.110	1.55	0.98	0.65	7	2.050
tau Bootis	F6IV	1.30	3.313	0.046	-	1.02	0.13	1	0.044
70 Virgo	G4V	0.90	116.600	0.430	4.48	1.33	0.56	3	0.512
HD 114762	F9V	1.20	84.050	0.300	3.23	0.96	0.40	3	0.369
HD 110833	K3V	0.73	270.040	0.800	1.30	0.66	0.38	5	0.712
BD-04 782	K5V	0.67	240.920	0.700	1.16	0.59	0.34	5	0.635
HD 112758	K0V	0.79	103.220	0.350	1.17	0.50	0.25	4	0.340
HD 98230	F8.5V	1.30	3.980	0.060	-	1.22	0.15	1	0.052
HD 18445	K2V	0.73	554.670	0.900	1.36	0.79	0.50	6	1.219
HD 29587	G2V	0.98	1157.843	2.500	1.65	1.04	0.69	7	2.180
HD 140913	G0V	1.05	147.940	0.540	1.68	0.71	0.36	4	0.488
HD 283750	K2	0.73	1.790	0.040	-	0.55	0.07	1	0.024
HD 217580	K4V	0.70	454.660	1.000	1.12	0.65	0.41	6	0.999
Alpha Tau	K5III	1.20	654.000	1.350	1.61	0.93	0.59	6	1.437
Prox.Cent.		0.10	42		0.20	0.10	0.06	5	0.111
Barnard's		0.12	132.0		0.19	0.12	-	7	0.249

V. Quantized galactic redshifts

Starting from the seventies, various authors ([?], [?], [?],[?], [?]) found that the frequency analysis of the galactic redshifts (both for binary galaxies and in the general case) indicates speed differences which are multiples of 37.5 km/s (Tifft suggests 36 km/s). Using our model, one can try to explain both this discretization and the estimated speed periodical structure as effects of the speed quantization rule

$$v_n = \frac{\alpha_g c}{n} = \frac{143.7}{n} \text{ km/s} .$$

VI. Spin of the celestial bodies

In order to validate or disprove the effectiveness of our scheme and the universality of the calculated α_g , we have taken consideration the known empirical the relationship between the angular momentum J of a celestial body and its mass M . Studying this question, both from a theoretical and a phenomenological point of view, Brosche (see for instance [?], [?], [?]) and later Wesson ([?], [?]) suggested that the law

$$J = pM^2$$

holds for purely gravitationally bounded systems. Wesson, fitting the empirical values and taking into account some theoretical consistency constraints suggested to use the value

$$p \simeq 8 \cdot 10^{-16} \text{ cm}^2 \text{ g}^{-1} \text{ s}^{-1}.$$

Consistently with the correspondence $e^2 \longleftrightarrow GMm$ on which our mechanical model is based, and recalling that the electron spin can be written in the form

$$s = \frac{1}{2} \frac{e^2}{\alpha_e c},$$

we can suppose that the rotational momentum of a purely gravitationally bound celestial body is given by

$$J = \frac{1}{2} \frac{GM^2}{\alpha_g c}.$$

As a consequence, we get $p = G/(2\alpha_g c) = 2.32 \cdot 10^{-15} \text{ cm}^2 \text{ g}^{-1} \text{ s}^{-1}$.

We note that, independently from this new guess, the previous formula can also be directly obtained from our considerations on the discretization of the orbital motions if a flat continuous purely gravitating disk is considered in which each infinitesimal mass ring rotates on the Bohr radius of the internal part of the disk which it delimits. In this case, the total angular momentum can be easily calculated by the formula

$$J = \int dJ = \frac{G}{\alpha_g c} \int m \, dm,$$

which just gives the previously guessed expression. In fact, this remark suggests that the concerned expression works well for flat or almost flat celestial

bodies (prevailing gravitational energy) whereas some correction factor can be needed in the opposite cases (when both the gravitational and the electrical energy play their role).

Anyway, we have compared these theoretical spin values with empirical values found in the literature or, as in the case of most planets, with the values

$$J_{obs} = \frac{2}{5}MR^2\frac{2\pi}{P_{obs}},$$

where P_{obs} indicates the observed rotation period (note that this formula refers to spherical and homogeneous rigid bodies and in fact it gives upper limits for the actual values).

In Fig.2 we report the logarithmic graph of our expression $J = pM^2$ compared with the experimental data: since no fit was performed, the agreement between our formula and the data (over more than twenty orders of mass magnitude) can be judged quite good.

In fact, the comparison shows some small discrepancy for the the solar planets, which on the other hand are examples of celestial bodies held together by other forces besides the gravitational one.

In the following table the calculated and the "observed" (upper limits) angular momenta for some planets are shown: the Earth apart, only for Saturn, owing to its oblate shape, the angular momentum $J = 8.16 \cdot 10^{37} Js$ can be directly calculated without using the previously remarked assumptions about the spherical planet form and its mass distribution.

A clear discrepancy regards the Earth, which shows $J_{calc} = 8.24 \cdot 10^{33} Js$ to be compared with $J_{obs} = 5.88 \cdot 10^{33} Js$. This fact can probably be accounted by the permanent loss of angular momentum caused by the tidal friction. An increasing rate of the rotation period of the Earth of about $16 s/10^6 years \simeq 3 \cdot 10^{-12}$ has been estimated [?]. This would correspond to a Moon braking action beginning 'only' about 1600 thousand millions of years ago, long after the Earth ocean was formed. On the other hand, the discrepancy in question between calculated and measured angular momentum of the Earth would be even greater if the whole Earth-Moon system ($J_{obs} = 3.47 \cdot 10^{34} Js$) was considered. Both this and the previous remarks seem to indicate that the Moon and the Earth can not be considered as a single celestial body: both could have originated from protoplanetary material on the same orbital ring ($n = 5$, $a_5 \simeq 1.07 a.u.$) and subsequently have gravitationally captured each the other (actually the Earth captured the Moon) about $2 \cdot 10^9 years$ ago.

This hypothesis would also account for the 9% of discrepancy between the observed and calculated terrestrial orbital radius round the Sun.

Body	J_{calc} [$J\ s$]	J_{obs} [$J\ s$]
Mercury	$2.53\ 10^{31}$	$< 1.02\ 10^{30}$
Venus	$5.50\ 10^{33}$	$< 2.14\ 10^{31}$
Earth	$8.24\ 10^{33}$	$5.88\ 10^{33}$
Mars	$9.56\ 10^{31}$	$< 2.10\ 10^{32}$
Jupiter	$8.37\ 10^{38}$	$< 6.83\ 10^{38}$
Saturn	$7.50\ 10^{37}$	$8.10\ 10^{37}$
Uranus	$1.76\ 10^{36}$	$< 2.50\ 10^{36}$
Neptune	$2.46\ 10^{36}$	$< 2.30\ 10^{36}$
Pluto	$5.23\ 10^{28}$	$< 1.5\ 10^{29}$

VII. Conclusions

- Assuming that the previously stated analogies hold between celestial and Old Quantum Mechanics, one can calculate values for a number of celestial quantities which turn out to be in remarkable agreement with the corresponding observed values.
- The gravitational structure constant $\alpha_g \simeq 1/2086$ has been used in order to account some observed values related to
 - the planetary motions around the Sun
 - the structure of the Saturn rings
 - the extra-solar planetary systems
 - the quantization of the galactic red-shifts
 - the spin of a broad class of celestial bodies, from solar planets to supercluster.
- No definite theory has yet been devised on the why the concerned analogies hold. Nevertheless, Nelson's stochastic mechanics ([?], [?]) seems at present the best candidate to offer valuable suggestions. In fact, in Nelson's scheme, a particle of mass which moves in a potential field is always subjected to a Brownian motion with diffusion coefficient

$\hbar/2m$: thus, the particle motion can be described using a probability density which turns out to obey the Schrödinger equation (of which the rules of the Old Quantum mechanics are consequences). If one consider the gravitational potential of a source of mass M , the diffusion coefficient of the appropriate Brownian motion would be $GM/(2\alpha_g c)$. This approach has been already applied to the study of the solar system, with some interesting results ([?], [?]).

In fact, we suspect that the Nelson's stochastic approach is an effective way to describe a purely classical deterministic chaotic dynamics, where the chaotic behaviour arises from the strong nonlinearities in a many body problem. The verified existence of stable attracting orbits in several many body problems can be a clue to the truth of this hypothesis.

A different interesting approach, which gives results very similar to those previously mentioned, has been worked out by Nottale (see for instance [?],[?]). It is based on a scale relativistic scheme, where the space-time resolution is an essential physical variable, and the physical laws are to be scale-covariants under resolution transformations: the fractal structure of the space-time plays a rôle of capital importance in the theory.

Acknowledgments

We are much obliged to H. Arp for his kind encouragement and his interest in our work, and for some important piece of information he supplied us about the work of both L. Nottale and N. Oliveira. We also express our thanks to P. Wesson for his interesting suggestions and his pleasant call. One of us thanks E.I. Guendelman for the kind hospitality in Beer Sheva.